

To Request More Information:

Please answer the following questions and mail to Deb at the address below.

I am a: FOD Family FOD Familymember Professional
Check all that apply: /my child has an FOD Please add me to your Mailing List Only

Comments:

Name and/or Professional Title

Address 1

Address 2

Phone [Deb calls new Families in the US and abroad]

Fax

Email

Please mail to:

**FOD Family Support Group
Deb Lee Gould, MEd, Director
PO Box 54
Okemos, MI 48805**

The FOD Family Support Group is an all volunteer family-based Support Group and a 501c3 tax-exempt Non-Profit Corporation. **Donations are tax deductible.** Your donation will help us cover copying, postage, website fees, phone calls to US and abroad, and Conference expenses, as well as for Clinical/Research Funds. No donations are used for administrative salaries. If you write a US check, please make it out to the 'FOD Family Support Group' and mail to the address above. We also have a secure PayPal donation link on our homepage. You can designate your donation to the General FOD Fund (daily operations), FOD Clinical Fund, FOD Research Fund or Grief Consults. **THANK YOU from ALL of our Families!**

For Families Living with a Fatty Oxidation Disorder



Fatty Oxidation Disorders Family Support Group

www.fodsupport.org



Deb Lee Gould, MEd, Director
MCAD Parent and Grief Consultant
PO Box 54
Okemos, MI 48805

Phone: 517.381.1940
Fax: 866.290.5206

Email: deb@fodsupport.org
Website: www.fodsupport.org

501c3 tax-exempt Non-Profit Corporation
Federal Tax ID # 83-0471342

FOD Family Support Group

Our History and Mission

The Fatty Oxidation Disorders (FOD) Family Support Group was 'born' in 1991 as a way of dealing with the sudden death of our 21-month-old daughter, Kristen, in 1985 from undiagnosed MCAD. Initially she was 'diagnosed' with Reye's Syndrome and it was not until one year later, when our son, Kevin, was born and diagnosed with MCAD that we discovered Kristen also had this rare metabolic disorder. Our third child, Brian, is a carrier and not directly affected. To say the least, our Family has been changed forever by this genetic disorder.



*Our hearts
give immortality to those
we love in memory...*

Our Mission is clear...to connect and network with FOD Families and Professionals around the world. Through our online newsletters, website, phone calls, and Email List, we provide emotional support, practical information about living with these disorders, and inform Families of new developments in screening, diagnosis, treatment, and research.

Awareness of these disorders is imperative for early diagnosis and treatment, prompting us to be committed advocates for Expanded Universal and Comprehensive Newborn Screening and longterm follow-up care for FODs and several other metabolic disorders. We do not want other Families to needlessly go through what we experienced.

From our Family to yours, we want you to know...

**You are NOT alone
and
'We Are All in This Together'**

**Take care...
Deb and Dan
Kristen, Kevin and Brian Gould**



FODs Defined

Fatty Oxidation Disorders are genetic metabolic deficiencies in which the body is unable to oxidize (breakdown) fatty acids to make energy because an enzyme is either missing or not working correctly. The main source of energy for the body is a sugar called glucose. Normally when the glucose runs out, fat is broken down into energy. However, that energy is not readily available to a child or adult with an FOD. If undiagnosed and untreated, these disorders can lead to serious complications affecting the liver, heart, eyes, and general muscle development, and possibly death. These disorders are sometimes misdiagnosed as SIDS and Reye's Syndrome.

Symptoms

There is a wide variation in the presentation of FODs, even within the same Family. Not every individual responds to a disorder in the same way. Some may be without major symptoms, yet others may have chronic bouts of low blood sugar or illnesses leading to hospitalizations. Please note that **blood sugar levels should not be totally relied on as the only indicator of a possible crisis** - the levels may even look normal. However, some of our children/adults are symptomatic with glucose levels of even 70 or 80mg/dl. Do NOT compare FODs with diabetes! Changes in behavior, irritability, lethargy (sleepy), and blood levels such as liver enzymes, acylcarnitines, ammonia, and ck (cpk) should also be monitored.

An emergency situation exists when a 'metabolic crisis' occurs. A crisis is often preceded by a period of fasting, possibly due to vomiting or an infection. Other symptoms may include diarrhea, seizures, coma and difficulty breathing.

Awareness of how to diagnose and treat these disorders is vital, because during a metabolic crisis, an undiagnosed individual may experience excessive buildup of fat in the liver, heart and kidneys, along with some brain swelling...all of which can lead to death.

Diagnosis and Treatment

Fatty Oxidation Disorders are autosomal disorders affecting both males and females. Both parents have to be at least a carrier of an abnormal gene (we do have parents with the disorder as well), in that when two abnormal genes unite the child will have an FOD. There is a 25% chance that EACH child will have an FOD and a 75% chance of being either a carrier or not having the disorder at all. If one child is diagnosed with an FOD, their siblings should also be tested, even if they are asymptomatic.

These disorders are screened for and/or diagnosed in a variety of ways at specialized Labs across the country (refer to our website). Blood, urine, skin fibroblasts, amniocytes (from amniotic fluid), and muscle and liver tissue are some of the specimens analyzed. The diagnostic tests often include an acylcarnitine profile, urine organic acid analysis, carnitine levels, and enzyme assays in fibroblasts. The acylcarnitine profile with whole blood on a 'PKU card' (filtered paper card) is the most direct approach for diagnosis of most of the FODs. Molecular DNA testing is also available for some FODs when the profile is abnormal. The Expanded Newborn Screen is a different test than the diagnostic profile. The bloodspot screen can 'red flag' 50+ different disorders, including many FODs.

One CAN live with an FOD! There may be various challenges along the way for some individuals with FODs (i.e., especially if diagnosed late, after a severe crisis, etc) but **KNOWING** you have an FOD and **how to treat** on a daily basis and in an emergency is the **Key!**

Treatment for FODs is multidimensional. A major concern is to avoid fasting. It is strongly recommended that infants under 6 months be fed around the clock every 3-4 hours. Older children or adults with an FOD eat often throughout their waking hours (i.e., every 3-6 hrs) and can sleep 10-12 hrs, when well. The feeding interval may depend on various factors and their specific circumstances, such as if they are ill versus well, in which case they most likely would eat more often around the clock. Please note that eating timeframes can vary from person to person within the different disorders.



Early Screening, Diagnosis and Treatment will prevent needless deaths

Disorders of Fatty acid Metabolism

Medium Chain acyl coA Dehydrogenase Deficiency (MCAD), Medium Chain 3-Ketoacyl coA Thiolase Deficiency (MCKAT), Medium/Short Chain L-3 Hydroxy acyl coA Dehydrogenase Deficiency (M/SCHAD), Long Chain L-3 Hydroxy acyl coA Dehydrogenase Deficiency (LCHAD), Very Long Chain acyl coA Dehydrogenase Deficiency (VLCAD), Short Chain acyl coA Dehydrogenase Deficiency (SCAD), 3-Hydroxy acyl coA Dehydrogenase Deficiency (HADH, formerly called SCHAD), Short Chain 3-Ketoacyl coA Thiolase Deficiency (SKAT, 3-ketothiolase), Electron Transfer Flavoprotein or Electron Transfer Flavoprotein ubiquinone-oxidoreductase Deficiency (ETF and ETF-QO, also known as GALL or MADD), Carnitine Palmitoyltransferase Deficiency (CPT I & II), Trifunctional Protein Deficiency (TFP), Carnitine Acylcarnitine Translocase Deficiency (CACT), Carnitine Transport Defect (Primary Carnitine Deficiency), 2,4-Dienoyl coA Reductase Deficiency, 3-Hydroxy 3-Methylglutaryl Deficiency (HMG), Unclassified FODs

A fasting state, especially while ill, can trigger a 'metabolic crisis' leading to lethargy and hospitalization. If hospitalized, it is imperative according to FOD experts, that a 10% dextrose IV is started immediately following blood chemistry samplings. Waiting hours for the lab test results before putting in the IV can be fatal when an FOD child/adult is in crisis.

[**Note:** Experts also recommend avoiding aspirin, oral products that contain salicylates, long term steroid medications, lactated ringers, and fat binding anesthesia medication].

Several snacks and meals of lowfat/high carbohydrate foods such as pasta, cereal, and other high complex carbohydrates (as well as sugar drinks when not well) are recommended throughout the day. Additionally, some Families use MCT Oil (not used for all FODs - i.e., MCAD and SCAD, since the medium chain triglycerides cannot be broken down). Hydrate with caloric drinks such as Gatorade while in heat and during exercise. Infants under 6 months should continue to have at least one night feed/snack as they should not go 10-12 hours without eating. If the FOD child/adult has experienced hypoglycemia (even with frequent meals), in order to help decrease the frequency of low blood sugar in the morning, some find it helpful to mix in 1-3 tablespoons of raw cornstarch (i.e., Argo) to a cold sugar-free liquid or food at night. Be aware that too much cornstarch can be constipating for some and may be hard on teeth. Be sure to brush after eating or drinking it. Also note that cornstarch should not be given to an infant under age one (they lack the necessary pancreatic amylase to digest the cornstarch).

Some physicians also may prescribe an L-carnitine drug supplement (FDA-approved Carnitor® or an FDA-approved generic form—these are NOT the same as the healthfood store carnitine), especially if a secondary carnitine deficiency is involved. Although carnitine is made within the body, sometimes supplementation is necessary. It is useful in stabilizing blood sugar levels, increasing muscle tone, and removing metabolic waste products from the cells so a buildup of toxins will not occur.

As with any medical condition, discussing treatment options with your physician(s) and FOD specialists is necessary for the well-being of anyone with an FOD. Please do **NOT** make any changes in your child's or your own treatment without first consulting with your physicians.

Additional Metabolic Family Support Resources

Organic Acidemia Association (OAA)

Kathy Stagni, Director
13210 35th Avenue North
Plymouth, MN 55441
763.559.1797
www.oaanews.org
kstagni@oanews.org

United Mitochondrial Disease Foundation (UMDF)

8085 Saltzburg Road, Suite 201
Pittsburgh, PA 15239
412.793.8077 or toll free 888.317.8633
412.793.6477 fax
www.umdf.org
info@umdf.org

Children Living with Inherited Metabolic Disease (CLIMB)

Climb Building
176 Nantwich Road
Crewe CW2 6BG
United Kingdom
0800 652 3181
0845 241 2174 fax
www.climb.org.uk
Info.svcs@climb.org.uk

Expanded NBS & FOD Diagnostic Testing:

[more Labs are listed on www.fodsupport.org, NBS/Medical Info page]

Mayo Medical Laboratories (Rochester, MN)

Mayo Clinic Biochemical Genetics Laboratory
Phone: 800.533.1710 for NBS testing,
Diagnostic testing, Consultation, Post-mortem testing
Fax: 507.284.1759
www.mayoclinic.org/laboratorygenetics-rst/biochemical.html

National Newborn Screening and Genetics Resource Center (NNSGRC) (Austin, TX)

512.454.6419
<http://genes-r-us.uthscsa.edu/index.htm>

Sigma-Tau Pharmaceuticals, Inc.

Makers of Carnitor®

9841 Washingtonian Blvd, Suite 500
Gaithersburg, MD 20878
800.447.0169
www.sigmtau.com
sigmtauinfo@sigmtau.com

FOD Family Support Group Board of Directors

Deb Lee Gould, MEd
FOD Parent, Director, and President

Daniel R Gould, PhD
FOD Parent and Treasurer

Mary Lingle
FOD Parent, Webmaster, and Secretary

Honorary Directors

Kevin Gould, MCAD Adult
Brian Gould, MCAD Sibling

Board of Medical Advisors

Charles R Roe, MD
Consultant
Dallas, TX

Mark Korson, MD
Tufts-New England Medical Center
Boston, MA

Jerry Vockley, MD, PhD
Children's Hospital of Pittsburgh
Pittsburgh, PA

Stephen G Kahler, MD
University of Arkansas for Medical Sciences
Little Rock, AR

Gary Siskin, MD
Albany Medical College
Albany, NY

Andrew Morris, PhD, FRCPC
Royal Manchester Children's Hospital
Manchester, UK

Arnold Strauss, MD
Cincinnati Children's Research Foundation
Cincinnati, OH

Barb Marriage, PhD, RD
Ross Metabolics/Abbott Nutrition
Columbus, OH

Janice Fletcher, MD, FRACP, FRCPA
Women and Children's Hospital
North Adelaide, Australia

James Gibson, MD, PhD
Specially for Children
Austin, TX



You are not alone...